

Synthesis, Characterization and Evaluation of Some Green Mannich Nano-Dispersant from Egg-Albumin

Rabab M. Nasser*^{1, 2}, Asmaa G. Komier², Bushra Y. Madkhali²

¹ Petroleum Applications Department, Egyptian Petroleum Research Institute, Nasr City, Cairo, Egypt

² Chemistry Department, University College of Samtah, Jazan University, Saudi Arabia

Received April 29, 2020; Accepted August 6, 2020


Abstract

The reaction of various aldehydes (formaldehyde, benzaldehyde, acetaldehyde and salicylaldehyde), albumin as an amine, and benzophenone as a ketone were synthesized into the four green basements of Mannich. The prepared compounds were characterized by determination of melting point and the structure was elucidated using FTIR spectrophotometer. The particle size, the zeta potential parameters and the morphological properties for the prepared compounds were characterized. The effect of the prepared compounds as oil spill dispersants for sweet water was evaluated and they have good efficiency as oil spill dispersants.

Keywords: Mannich base; Egg – albumin; FT-IR; Zeta potential; SEM, Oil spill dispersion.

1. Introduction

The end product of Mannich reaction is the Mannich base, beta-amino ketones containing compounds [1-4]. The Mannich reaction is a nucleophilic replacement reaction involving an amino (primary or secondary) and formaldehyde (any Aldehyde) compound condensation with active hydrogen [5]. The general Mannich reaction is represented in Scheme 1.



Scheme 1. Mannich base formation

Studies have shown Mannich bases to be highly reactive and to be convertible easily into compounds such as reduced to physiologically active amino alcohols [6]. It has been stated that Mannich bases have powerful action, such as anti-inflammatory activity [7-8], anticancer [9-10], anti-filarial activity [11]. Mannich bases are well known for their use in detergents additives [12-13], resins, polymers, surface active ingredients [14] and so on as well as in biological activity. Different active compounds have been able to solve these constraints with Mannich drugs [15]. For the synthesis of bioactive molecules, the Mannich bases and their derivatives are intermediates [16-17]. The Mannich reaction is widely used in nitrogen compound construction. Because of their use in antibacterial action [18], Mannich bases gained popularity and other applications are in agrochemical products such as plant growth regulators. In the present work four green Mannich bases were synthesized and characterized by determination of melting point, FT-IR spectrophotometer, zeta potential and the morphological structure was carried out by scanning electron microscope. The effect of the prepared compounds as oil spill dispersants was studied.

2. Materials and methods

2.1. Materials

Formaldehyde; LOBA CHEMIE 99.00%, acetaldehyde; JANSSEN CHIMICA 99.00%, benzaldehyde; LOBA CHEMIE 100%, salicylaldehyde; HIMEDIA 99.00%, benzophenone; LOBA CHEMIE 99.00% and albumin; LOBA CHEMIE 99.50% were used without further purification.

2.2. Methods

Four Mannich bases were synthesized using (2:1:2) molar ratios of aldehydes (formaldehyde, acetaldehyde, benzaldehyde and salicylaldehyde) separately, ketone (benzophenone) and amine (egg- albumin). The reaction was carried out at 70-100°C, using 1 mL of concentrated hydrochloric acid as a catalyst for water removal.

2.2.1. Recrystallization process

The recrystallization process was carried out. The prepared compounds were recrystallized by heating with hot water as a solvent for recrystallization, after complete dissolving of the solute; the crystallization beaker was removed and cooled at room temperature. The beaker was then put in an ice-water bath and left to form crystals. Using vacuum filters, the crystals were collected and dried.

2.2.2. Determination of melting points for the prepared compounds

The melting points of the prepared compounds were measured using STUART SMP10.

2.2.3. Characterization of compounds prepared with FT-IR.

The prepared compounds were characterized using Fourier Transformer Infra- Red Spectroscopy, Model Nicolet iS10-FTIR Spectrophotometer, KBr.

2.2.4. Zeta parameters characterization of the compounds prepared

The parameters of the prepared compounds were characterized by laser Doppler anemometry on a Malvern zeta-sizer, dynamic light scattering with a (3000 HSA, Malvern Instruments, UK) at 90°C at a scattering angle. This was done via the average particle size, size distribution and multipoint indicators. Dilution of formulated compounds suspensions with deionized water was used to obtain specimens. The average value \pm standard deviation (S.D.) was stated per value for three measurements, Table 2.

2.2.5. The morphological characteristics of prepared compounds

Scanning electron microscope – QUANTA- FEG 250 was used for the microscopic characterization of the prepared compounds. The energy used for the acceleration beam was 20 KV. The magnification power (40-500 μ m) of all the micrograms was taken.

2.2.6. Assessment of the prepared compounds as oil spill dispersants.

The prepared compounds were evaluated as oil spill dispersants for palm oil according to the following methods:

2.2.6.1. SFT Oil extraction and analysis

The determination of the efficiency index of oil spill dispersants was made in the test flask, followed by the sequence addition of oils, and finally the dispersant, with a volume of 120mL of tap-water equilibrated to the desired temperature. Subsequently, the 5 mL syringe tip attachment on the water surface instantly dispensed with 100 mL of oil. The dispersant was then distributed in the center of the slick which was fitted with a 1:25 Dispersant – to – oil relation (DOR) with a 100mL syringe tip fixture up to 4 mL. The flask was put in an orbital shaker for 10 minutes, the shaker was maintained at the desired rotating velocity at the end, and it was removed. At the end of the setting period, the first 2 ml sample was dried and removed and a 30 mL sample was taken in 50 mL cylinder to a separating funnel with 125 mL.

New 5 mL of three extracts using dichloromethane is made. The extract must then be modified to a final volume of 20 mL and transferred into a 50 mL crimp bottle with a teflon/aluminum seal. The glasses are maintained at $4 \pm 2^\circ\text{C}$ (maximum 5 days) before examination. The methodology developed is accompanied by the oil standard process and the test procedures [19].


2.2.6.2. Efficiency of the prepared compounds as oil spill dispersants

The efficiency of the effect of the four prepared compounds according to (IP-AS/84) standard method at 20°C . The efficiency index (E) was calculated according to the following equation:

$$E = \frac{\text{Weight of oil in } 50\text{cm}^3 \text{ sample of oily water} \times 500}{\text{Total weight of oil added to } 250\text{cm}^3 \text{ separating funnel}}$$


3. Results

Four green Mannich bases were synthesized by reaction of different aldehydes (formaldehyde, acetaldehyde, benzaldehyde, and salicylaldehyde) separately with egg albumin and (benzophenone) as a ketone. The feed ratio was 2:1:2 molar, Schemes (2-5).




N,N-bis((4-(hydroxy(phenyl)methylene)cyclohexa-2,5-dien-1-yl)methyl)-5,5-dimethyl-2,4-dioxooxazolidine-3-carboxamide

Scheme 2. A₁ Mannich base.




N,N-bis(1-(4-(hydroxy(phenyl)methylene)cyclohexa-2,5-dien-1-yl)ethyl)-5,5-dimethyl-2,4-dioxooxazolidine-3-carboxamide

Scheme 3. A₂ Mannich base



Scheme 4. A₃ Mannich Base.



Scheme 5. A₄ Mannich Base.

The prepared compounds (A₁-A₄) were recrystallized and melting points were determined, data is tabulated in Table 1. The yield% of the prepared compounds was calculated according to the following equation:

$$\text{Yield \%} = \frac{\text{actual yield}}{\text{theoretical yield}} \times 100$$

The data was tabulated in Table 1. It was obvious that the A₄ compound has the greatest Yield %. This may be due to A₄ compound has the highest molecular weight of the prepared compounds.

Table 1. The designation, chemical properties of (A₁-A₄) compounds


Cpd.	Cpd. designation	M.p.(°C)	Yield %	Molecular formula	M. wt.
A ₁	Formaldehyde: Albumin: benzo-phenone	146-148	92	C ₃₄ H ₃₂ N ₂ O ₆	564.63
A ₂	Acetaldehyde: Albumin: benzophe-none	198-200	93	C ₃₆ H ₃₆ N ₂ O ₆	592.68
A ₃	Benzaldehyde: Albumin: benzo-phenone	192-194	92	C ₄₆ H ₄₀ N ₂ O ₆	716.82
A ₄	Salicyaldehyde: Albumin: benzo-phenone	184-186	95	C ₄₆ H ₄₀ N ₂ O ₈	748.82

The zeta potential parameters were determined as tabulated in Table 2.

Table 2. Zeta potential parameters of the prepared compounds

Sam-ple	Dispersant name	RI	Viscosity (cP)	DDC	Conductivity (ms/cm)	ZP (mv)	ZD (mv)	Average size - z (d. nm)	pdi
ALB.					0.209	-13.8	-13.8±4.48	401.7	0.731
A ₁	water	1.33	0.8872	78.5	0.0141	-4.39	-4.39±4.38	1386	0.895
A ₂					0.726	5.96	5.96±4.65	4418	0.321
A ₃					0.178	8.08	8.08±3.66	2402	1.000
A ₄					0.153	-0.0857	-0.0857±4.74	1870	0.384

The prepared Mannich bases were subjected to surface morphology study by scanning electron microscopy QUANTA- FEG 250, using liquid nitrogen, Figure 1.

Figure 1. SEM morphology for compound A₁

4. Discussions

The chemical structure of the prepared compounds was described with FT-IR spectroscopy, from FT-IR spectra, Figure 2. By comparing FT-IR spectra of the prepared compounds with the FT-IR spectra of albumin one can concluded that the disappearance of the –NH₂ peaks at 3442 and 3360cm⁻¹. This indicate the completely conversion of –NH₂ group into tertiary amine group.




Figure 2. FT-IR of albumin and the prepared (A₁-A₄) compounds.

The specific physical and chemical properties of palm oil were characterized; the data were tabulated at Table 3.

Table 3. Physicochemical properties of palm oil

Physicochemical properties	Palm oil	SON standard
Smoke point (°C)	121	--
Flash point (°C)	182	--
Fire point (°C)	196	--
Moisture (%)	0.24	0.29
Specific gravity	0.824	0.897-0.907
Free fatty acids	3.10	3.5
Iodine value	29.56	45-53
Peroxide value	7.90	10
Saponification value	195.20	195-205

To obtain the best results the dispersant used has to be carefully selected. The effect of any dispersant depends on many factors, such as the nature of the dispersant itself, the nature of the oil spilled, and the temperature [20-21]. Thus, dispersants efficiencies on the oil were determined and given in Table 4. The results obtained show that in case of chemical dispersants (A₁-A₃), having two OH group, accordingly they have good dispersing effect ranging from (63 -74), while compound (A₄) has the highest dispersing power (88), this may be due to the presence of four OH groups besides the increase of the electron cloud inside the molecule.

Table 4. Efficiency of dispersants on oil at 20°C

Dispersant	Weight of oil in 50cm ³ sample of oily water	Total weight of oil added to 250cm ³ separating funnel	Efficiency @ 20°C
A ₁	1.26	10	63
A ₂	1.30		65
A ₃	1.48		74
A ₄	1.76		88

Different ratios were taken to determine the relation between the efficiency and DOR. As it obvious from Figure 3, it was found that the highest efficiency of the used compounds (A₁-A₄) according to DOR method is 1:50.




Figure 3. Efficiency of (A₁-A₄) on palm oil

5. Conclusions

In the present work four Mannich bases were synthesized and characterized by determination of melting point. The yield % of the prepared compounds was calculated and it was found that, A₁ compound has the highest yield. The chemical structure of the prepared compounds was elucidated using FT-IR spectroscopy. The FT-IR spectra of the prepared compounds illustrate the disappearance of -NH₂ group of albumin and indicate the complete conversion of it into tertiary amine. Zeta parameters were characterized. The prepared compounds were evaluated as oil spill dispersants and they were obtained a good oil spill dispersant effect.

Abbreviations

FT-IR	Fourier Transformer Infra-Red spectrophotometer;
DOR	dispersant oil ratio;
SFT	the swirling flask test
E	efficiency index
RI	refractive index
DDC	dispersant dielectric constant
ZP	Zeta potential
ZD	Zeta deviation
pdi	polydispersity index

Acknowledgement

The research was financially supported by Deanship of Scientific Research, Jazan University, Saudi Arabia. Also the authors would like to thank Dr. S.M. Siva Kumar, for helping in zeta analysis.

Author contributions

The manuscript was written through contributions of all authors. Rabab M. Nasser, suggested the idea and contributed the writing of the paper. Asmaa G. Komier and Bushra Y. Madkhali help in the experimental part of the work.

Funding sources

This project was supported by Jazan University, Deanship of Scientific Research.

References

- [1] Idhayadhulla A, Surendra RK, Nasser AJA, Selvin J. Synthesis of some Mannich base derivatives and their antimicrobial activity study. *Arabian Journal of Chemistry*, 2014; 7: 994–999.
- [2] Idhayadhulla A, Kumar R S, Nasser AJA, and Manilal A. Synthesis and antimicrobial activity of some new Mannich base derivatives. *Journal of Chemical and Pharmaceutical Research*, 2011; 3(4): 904–911.
- [3] Belinelo VJ, Reis GT, Stefani GM, Ferreira-Alves DL, and Pilo-Veloso D. Synthesis of 6 α ,7 β -dihydroxyvouacapan- 17 β -oic acid derivatives. Part IV: Mannich base derivatives and its activities on the electrically stimulated guinea-pig ileum preparation. *Journal of the Brazilian Chemical Society*, 2002, 13(6): 830–837.
- [4] Joshi S, Khosla N, Tiwari P. In vitro study of some medicinally important Mannich bases derived from anti-tubercular agent. *Bioorganic & Medicinal Chemistry*, 2004; 12,(3): 571– 576.
- [5] Ayesha F, Muhammad I, Amna F, Shoomaila L, Muhammad L,, Zaigham A, Gabriel B, and Liviu M. antibacterial activity studies of Co(II), Ni(II), Cu(II) AND Zn(II) complexes with Mannich base ligand; *Bull. Chem. Soc. Ethiop.*, 2019; 33(3): 485-492.
- [6] Raman N, Esthar S, Thangaraja C. A new Mannich base and its transition metal (II) complexes—synthesis, structural characterization and electrochemical study. *Journal of Chemical Sciences*, 2004; 116(4): 209–213.
- [7] Hayun H, Arrahman A, Purwati EM, Yanuar A, Fortunata F, Suhargo F, Syafiqah DW, Ignacia C, Novalia AR. Synthesis, Anti-inflammatory and Antioxidant Activity of Mannich Bases of Dehydrozingerone Derivatives; *J. Young Pharm.*, 2018; 10(2): 6-10.
- [8] Meric K, Irem O-D, Tugce O, Beril K, Hande S, Ayhan B, and Suleyman SB. Some Novel Mannich Bases of 5-(3,4-Dichlorophenyl)-1,3,4-oxadiazole-2(3H)-one and Their Anti-Inflammatory Activity. *Arch. Pharm. Chem. Life Sci.*, 2017; 350: 1700153.
- [9] Bhupendra M, Rahul VP, Young SK, Rafi N, Enkhtaivan G, Kim DH. Synthesis of Mannich base derivatives of berberine and evaluation of their anticancer and antioxidant effects; *Journal of Chemical Research*, 2016; 40: 73–77.
- [10] Van-Son N, Ling Sh, Fang-Qian L, Qiu-An W. Synthesis of kaempferide Mannich base derivatives and their antiproliferative activity on three human cancer cell lines; *Acta Biochimica Polonica*; 2015; 62(3) :547–552.
- [11] Kalluraya, B.; Chimbalkar, R. M.; Hegde, J. C.; Anticonvulsant activity of nicotiny/isonicotinyl substituted 1, 2, 4- triazol-5-thione Mannich bases. *Indian Journal of Heterocyclic Chemistry*, 2005; 15(1): 15–18.
- [12] Nehal SA, Amal MN, Abd El-Aziz Kh, Azim A-AAA, El-Kafrawy AF. Deposit control agents for lubricating oil; *Petroleum and Coal*; 2017; 59(4): 453-463.
- [13] Nassar AM, Ahmed NS, Abd El-Aziz Kh., Azim A-AAA, El-Kafrawy AF. Synthesis and Evaluation of Detergent/Dispersant Additives from Polyisobutylene Succinimides; *International Journal of Polymeric Materials*, 2006; 55: 703–713.
- [14] Wang Y, Chu X, Zhang M. Tribological studies on a new borated mannich base containing benzotriazole group as additive for environmentally adapted lubricant; *Tribology*, 2016; 10(3): 95-100.
- [15] Li Y, Qiang X, Luo L, Yang X, Xiao G, Zheng Y, Cao Zh, Sang Zh, Su F, Deng Y. Multitarget drug design strategy against Alzheimer’s disease: Homoisoflavonoid Mannich base derivatives serve as acetylcholinesterase and monoamine oxidase B dual inhibitors with multifunctional properties; *Bioorganic & Medicinal Chemistry*, 2017; 25:714–726.

- [16] Zhang Y, Li Zh, Song H, Wang B. Structure and Biological Activities of Novel 2-(Trifluoromethyl) - 6-arylimidazo[2,1-b][1,3,4]-thiadiazole (bis-)Mannich Base Derivatives Containing Substituted piperazine Moiety. *J. Chem. Synthesis*, 2018; 36: 635–638.
- [17] Szatmári I, Belasri Kh, Heydenreich M, Koch A, Kleinpeter E, Fülöp F. Ortho-Quinone Methide Driven Synthesis of New O,N- or N,N-Heterocycles; *Chemistry Open* 2019; 8: 961–971.
- [18] Pandeya SN, Sriram D, Nath G, de Clercq E. Synthesis, Antibacterial, Antifungal and Anti- HIV Evaluation of Schiff and Mannich Bases of Isatin and its Derivatives with Triazole; *Arzneim.-Forsch. Drug Res.*, 2000; 50 (1): 55–59.
- [19] Sorial GA, Koran KM, Holder E, Venosa AD, and King DW. Development of a rational oil spill dispersant effectiveness protocol. *International Oil Spill Conference Proceedings*, 2001; 1: 471-478.
- [20] Garrett RM, Guénette ChC, Haith CE, and Prince RC. Pyrogenic Polycyclic Aromatic Hydrocarbons in Oil Burn Residues. *Environ. Sci. Technol.*, 2000; 34:1934–1937.
- [21] Damilola VA, Oyinkepreye D O, Vincent EE, Oluwasanmi O, Temiloluwa IO. The influence of surfactant concentration and surfactant type on the interfacial tension of heavy crude oil/ Brine/ Surfactant system. *Petroleum and Coal*, 2020; 62(2) 292-298.

To whom correspondence should be addressed: Dr. Rabab M. Nasser, Petroleum Applications Department, Egyptian Petroleum Research Institute, Nasr City, Cairo, Egypt, E-mail: rabab_nasser@yahoo.com